

PubMed ▼

Format: Abstract

Full text links



Neuropharmacology. 2016 Apr;103:16-26. doi: 10.1016/j.neuropharm.2015.12.017. Epub 2015 Dec 19.

Cannabidiol induces rapid-acting antidepressant-like effects and enhances cortical 5-HT/glutamate neurotransmission: role of 5-HT1A receptors.

Linge R¹, Jiménez-Sánchez L², Campa L², Pilar-Cuéllar F¹, Vidal R¹, Pazos A¹, Adell A³, Díaz Á⁴.

Author information

Abstract

Cannabidiol (CBD), the main non-psychotomimetic component of marijuana, exhibits anxiolytic-like properties in many behavioural tests, although its potential for treating major depression has been poorly explored. Moreover, the mechanism of action of CBD remains unclear. Herein, we have evaluated the effects of CBD following acute and chronic administration in the olfactory bulbectomy mouse model of depression (OBX), and investigated the underlying mechanism. For this purpose, we conducted behavioural (open field and sucrose preference tests) and neurochemical (microdialysis and autoradiography of 5-HT1A receptor functionality) studies following treatment with CBD. We also assayed the pharmacological antagonism of the effects of CBD to dissect out the mechanism of action. Our results demonstrate that CBD exerts fast and maintained antidepressant-like effects as evidenced by the reversal of the OBX-induced hyperactivity and anhedonia. In vivo microdialysis revealed that the administration of CBD significantly enhanced serotonin and glutamate levels in vmPFCx in a different manner depending on the emotional state and the duration of the treatment. The potentiating effect upon neurotransmitters levels occurring immediately after the first injection of CBD might underlie the fast antidepressant-like actions in OBX mice. Both antidepressant-like effect and enhanced cortical 5-HT/glutamate neurotransmission induced by CBD were prevented by 5-HT1A receptor blockade. Moreover, adaptive changes in pre- and post-synaptic 5-HT1A receptor functionality were also found after chronic CBD. In conclusion, our findings indicate that CBD could represent a novel fast antidepressant drug, via enhancing both serotonergic and glutamate cortical signalling through a 5-HT1A receptor-dependent mechanism.

KEYWORDS: (-)Cannabidiol (PubChem CID: 12302390); 5-HT(1A) receptor; AM251 (PubChem CID: 2125); Antidepressant; **Cannabidiol**; Glutamate; Olfactory bulbectomy; Serotonin; WAY-100635 maleate salt (PubChem CID: 11957721)

PMID: 26711860 DOI: [10.1016/j.neuropharm.2015.12.017](https://doi.org/10.1016/j.neuropharm.2015.12.017)

[Indexed for MEDLINE]

Publication type, MeSH terms, Substances



LinkOut - more resources



PubMed Commons

[PubMed Commons home](#)

0 comments

[How to join PubMed Commons](#)